

Credibility theory features of **actuar**

Christophe Dutang
Université Paris Dauphine

Vincent Goulet
Université Laval

Xavier Milhaud
Université Claude Bernard Lyon 1

Mathieu Pigeon
Université du Québec à Montréal

1 Introduction

Credibility models are actuarial tools to distribute premiums fairly among a heterogeneous group of policyholders (henceforth called *entities*). More generally, they can be seen as prediction methods applicable in any setting where repeated measures are made for subjects with different risk levels.

The credibility theory features of **actuar** consist of matrix `hachemeister` containing the famous data set of [Hachemeister \(1975\)](#) and function `cm` to fit hierarchical (including Bühlmann, Bühlmann-Straub), regression and linear Bayes credibility models. Furthermore, function `rcomphierarc` can simulate portfolios of data satisfying the assumptions of the aforementioned credibility models; see the "simulation" vignette for details.

2 Hachemeister data set

The data set of [Hachemeister \(1975\)](#) consists of private passenger bodily injury insurance average claim amounts, and the corresponding number of claims, for five U.S. states over 12 quarters between July 1970 and June 1973. The data set is included in the package in the form of a matrix with 5 rows and 25 columns.

The first column contains a state index, columns 2–13 contain the claim averages and columns 14–25 contain the claim numbers:

```
> data(hachemeister)
> hachemeister
```

	state	ratio.1	ratio.2	ratio.3	ratio.4	ratio.5
[1,]	1	1738	1642	1794	2051	2079
[2,]	2	1364	1408	1597	1444	1342
[3,]	3	1759	1685	1479	1763	1674
[4,]	4	1223	1146	1010	1257	1426
[5,]	5	1456	1499	1609	1741	1482

	ratio.6	ratio.7	ratio.8	ratio.9	ratio.10	ratio.11
[1,]	2234	2032	2035	2115	2262	2267
[2,]	1675	1470	1448	1464	1831	1612
[3,]	2103	1502	1622	1828	2155	2233
[4,]	1532	1953	1123	1343	1243	1762
[5,]	1572	1606	1735	1607	1573	1613

	ratio.12	weight.1	weight.2	weight.3	weight.4
[1,]	2517	7861	9251	8706	8575
[2,]	1471	1622	1742	1523	1515
[3,]	2059	1147	1357	1329	1204
[4,]	1306	407	396	348	341
[5,]	1690	2902	3172	3046	3068

	weight.5	weight.6	weight.7	weight.8	weight.9
[1,]	7917	8263	9456	8003	7365
[2,]	1622	1602	1964	1515	1527
[3,]	998	1077	1277	1218	896
[4,]	315	328	352	331	287
[5,]	2693	2910	3275	2697	2663

	weight.10	weight.11	weight.12
[1,]	7832	7849	9077
[2,]	1748	1654	1861
[3,]	1003	1108	1121
[4,]	384	321	342
[5,]	3017	3242	3425

3 Hierarchical credibility model

The linear model fitting function of R is `lm`. Since credibility models are very close in many respects to linear models, and since the credibility model fitting function of **actuar** borrows much of its interface from `lm`, we named the credibility function `cm`.

Function `cm` acts as a unified interface for all credibility models supported by the package. Currently, these are: the unidimensional models of [Bühlmann \(1969\)](#) and [Bühlmann and Straub \(1970\)](#); the hierarchical model of [Jewell \(1975\)](#) (of which the first two are special cases); the regression model of [Hachemeister \(1975\)](#), optionally with the intercept at the barycenter of time ([Bühlmann and Gisler, 2005](#), Section 8.4); linear Bayes models. The modular design of `cm` makes it easy to add new models if desired.

This section concentrates on usage of `cm` for hierarchical models.

There are some variations in the formulas of the hierarchical model in the literature. We compute the credibility premiums as given in [Bühlmann and Jewell \(1987\)](#) or [Bühlmann and Gisler \(2005\)](#), supporting three types of estimators of the between variance structure parameters: the unbiased estimators of [Bühlmann and Gisler \(2005\)](#) (the default), the slightly different version of [Ohlsson \(2005\)](#) and the iterative pseudo-estimators as found in [Goovaerts and Hoogstad \(1987\)](#) or [Goulet \(1998\)](#).

Consider an insurance portfolio where *entities* are classified into *cohorts*. In our terminology, this is a two-level hierarchical classification structure. The observations are claim amounts S_{ijt} , where index $i = 1, \dots, I$ identifies the cohort, index $j = 1, \dots, J_i$ identifies the entity within the cohort and index $t = 1, \dots, n_{ij}$ identifies the period (usually a year). To each data point corresponds a weight — or volume — w_{ijt} . Then, the best linear prediction for the next period outcome of a entity based on ratios $X_{ijt} = S_{ijt}/w_{ijt}$ is

$$\begin{aligned}\hat{\pi}_{ij} &= z_{ij}X_{ijw} + (1 - z_{ij})\hat{\pi}_i \\ \hat{\pi}_i &= z_iX_{izw} + (1 - z_i)m,\end{aligned}\tag{1}$$

with the credibility factors

$$\begin{aligned}z_{ij} &= \frac{w_{ij\Sigma}}{w_{ij\Sigma} + s^2/a}, & w_{ij\Sigma} &= \sum_{t=1}^{n_{ij}} w_{ijt} \\ z_i &= \frac{z_{i\Sigma}}{z_{i\Sigma} + a/b}, & z_{i\Sigma} &= \sum_{j=1}^{J_i} z_{ij}\end{aligned}$$

and the weighted averages

$$X_{ijw} = \sum_{t=1}^{n_{ij}} \frac{w_{ijt}}{w_{ij\Sigma}} X_{ijt}$$

$$X_{izw} = \sum_{j=1}^{J_i} \frac{z_{ij}}{z_{i\Sigma}} X_{ijw}.$$

The estimator of s^2 is

$$\hat{s}^2 = \frac{1}{\sum_{i=1}^I \sum_{j=1}^{J_i} (n_{ij} - 1)} \sum_{i=1}^I \sum_{j=1}^{J_i} \sum_{t=1}^{n_{ij}} w_{ijt} (X_{ijt} - X_{ijw})^2. \quad (2)$$

The three types of estimators for the variance components a and b are the following. First, let

$$A_i = \sum_{j=1}^{J_i} w_{ij\Sigma} (X_{ijw} - X_{iww})^2 - (J_i - 1)s^2 \quad c_i = w_{i\Sigma\Sigma} - \sum_{j=1}^{J_i} \frac{w_{ij\Sigma}^2}{w_{i\Sigma\Sigma}}$$

$$B = \sum_{i=1}^I z_{i\Sigma} (X_{izw} - \bar{X}_{zzw})^2 - (I - 1)a \quad d = z_{\Sigma\Sigma} - \sum_{i=1}^I \frac{z_{i\Sigma}^2}{z_{\Sigma\Sigma}},$$

with

$$\bar{X}_{zzw} = \sum_{i=1}^I \frac{z_{i\Sigma}}{z_{\Sigma\Sigma}} X_{izw}. \quad (3)$$

(Hence, $E[A_i] = c_i a$ and $E[B] = db$.) Then, the Bühlmann–Gisler estimators are

$$\hat{a} = \frac{1}{I} \sum_{i=1}^I \max\left(\frac{A_i}{c_i}, 0\right) \quad (4)$$

$$\hat{b} = \max\left(\frac{B}{d}, 0\right), \quad (5)$$

the Ohlsson estimators are

$$\hat{a}' = \frac{\sum_{i=1}^I A_i}{\sum_{i=1}^I c_i} \quad (6)$$

$$\hat{b}' = \frac{B}{d} \quad (7)$$

and the iterative (pseudo-)estimators are

$$\tilde{a} = \frac{1}{\sum_{i=1}^I (J_i - 1)} \sum_{i=1}^I \sum_{j=1}^{J_i} z_{ij} (X_{ijw} - X_{izw})^2 \quad (8)$$

$$\tilde{b} = \frac{1}{I - 1} \sum_{i=1}^I z_i (X_{izw} - X_{zzw})^2, \quad (9)$$

where

$$X_{zzw} = \sum_{i=1}^I \frac{z_i}{z_\Sigma} X_{izw}. \quad (10)$$

Note the difference between the two weighted averages (3) and (10). See [Belhadj et al. \(2009\)](#) for further discussion on this topic.

Finally, the estimator of the collective mean m is $\hat{m} = X_{zzw}$.

The credibility modeling function `cm` assumes that data is available in the format most practical applications would use, namely a rectangular array (matrix or data frame) with entity observations in the rows and with one or more classification index columns (numeric or character). One will recognize the output format of `rcomphierarc` and its summary methods.

Then, function `cm` works much the same as `lm`. It takes in argument: a formula of the form `~ terms` describing the hierarchical interactions in a data set; the data set containing the variables referenced in the formula; the names of the columns where the ratios and the weights are to be found in the data set. The latter should contain at least two nodes in each level and more than one period of experience for at least one entity. Missing values are represented by NAs. There can be entities with no experience (complete lines of NAs).

In order to give an easily reproducible example, we group states 1 and 3 of the Hachemeister data set into one cohort and states 2, 4 and 5 into another. This shows that data does not have to be sorted by level. The fitted model below uses the iterative estimators of the variance components.

```
> X <- cbind(cohort = c(1, 2, 1, 2, 2), hachemeister)
> fit <- cm(~cohort + cohort:state, data = X,
+          ratios = ratio.1:ratio.12,
+          weights = weight.1:weight.12,
+          method = "iterative")
> fit
Call:
cm(formula = ~cohort + cohort:state,
```

```
data = X, ratios = ratio.1:ratio.12,  
weights = weight.1:weight.12,  
method = "iterative")
```

Structure Parameters Estimators

Collective premium: 1746

Between cohort variance: 88981

Within cohort/Between state variance: 10952

Within state variance: 139120026

The function returns a fitted model object of class "cm" containing the estimators of the structure parameters. To compute the credibility premiums, one calls a method of predict for this class.

```
> predict(fit)  
$cohort  
[1] 1949 1543  
  
$state  
[1] 2048 1524 1875 1497 1585
```

One can also obtain a nicely formatted view of the most important results with a call to summary.

```
> summary(fit)  
Call:  
cm(formula = ~cohort + cohort:state,  
data = X, ratios = ratio.1:ratio.12,  
weights = weight.1:weight.12,  
method = "iterative")  
  
Structure Parameters Estimators  
  
Collective premium: 1746  
  
Between cohort variance: 88981  
Within cohort/Between state variance: 10952  
Within state variance: 139120026
```

Detailed premiums

Level: cohort

cohort	Indiv. mean	Weight	Cred. factor	Cred. premium
1	1967	1.407	0.9196	1949
2	1528	1.596	0.9284	1543

Level: state

cohort	state	Indiv. mean	Weight	Cred. factor
1	1	2061	100155	0.8874
2	2	1511	19895	0.6103
1	3	1806	13735	0.5195
2	4	1353	4152	0.2463
2	5	1600	36110	0.7398

Cred. premium

2048

1524

1875

1497

1585

The methods of predict and summary can both report for a subset of the levels by means of an argument levels.

```
> summary(fit, levels = "cohort")
```

Call:

```
cm(formula = ~cohort + cohort:state,  
    data = X, ratios = ratio.1:ratio.12,  
    weights = weight.1:weight.12,  
    method = "iterative")
```

Structure Parameters Estimators

Collective premium: 1746

Between cohort variance: 88981

Within cohort variance: 10952

Detailed premiums

```

      cohort Indiv. mean Weight Cred. factor Cred. premium
      1      1967      1.407  0.9196      1949
      2      1528      1.596  0.9284      1543
> predict(fit, levels = "cohort")
$cohort
[1] 1949 1543

```

4 Bühlmann and Bühlmann–Straub models

As mentioned above, the Bühlmann and Bühlmann–Straub models are simply one-level hierarchical models. In this case, the Bühlmann–Gisler and Ohlsson estimators of the between variance parameters are both identical to the usual [Bühlmann and Straub \(1970\)](#) estimator

$$\hat{a} = \frac{w_{\Sigma\Sigma}}{w_{\Sigma\Sigma}^2 - \sum_{i=1}^I w_{i\Sigma}^2} \left(\sum_{i=1}^I w_{i\Sigma} (X_{iw} - X_{ww})^2 - (I-1)\hat{s}^2 \right), \quad (11)$$

and the iterative estimator

$$\tilde{a} = \frac{1}{I-1} \sum_{i=1}^I z_i (X_{iw} - X_{zw})^2 \quad (12)$$

is better known as the Bichsel–Straub estimator.

To fit the Bühlmann model using `cm`, one simply does not specify any weights.

```

> cm(~state, hachemeister, ratios = ratio.1:ratio.12)
Call:
cm(formula = ~state, data = hachemeister,
    ratios = ratio.1:ratio.12)

Structure Parameters Estimators

Collective premium: 1671

Between state variance: 72310
Within state variance: 46040

```

When weights are specified together with a one-level model, `cm` automatically fits the Bühlmann–Straub model to the data. In the example below, we use the Bichsel–Straub estimator for the between variance.


```
> cm(~state, hachemeister, ratios = ratio.1:ratio.12,
+   weights = weight.1:weight.12)
```

Call:

```
cm(formula = ~state, data = hachemeister,
   ratios = ratio.1:ratio.12,
   weights = weight.1:weight.12)
```

Structure Parameters Estimators

Collective premium: 1684

Between state variance: 89639

Within state variance: 139120026

5 Regression model of Hachemeister

The credibility regression model of [Hachemeister \(1975\)](#) is a generalization of the Bühlmann–Straub model. If data shows a systematic trend, the latter model will typically under- or over-estimate the true premium of an entity. The idea of [Hachemeister](#) was to fit to the data a regression model where the parameters are a credibility weighted average of an entity's regression parameters and the group's parameters.

In order to use `cm` to fit a credibility regression model to a data set, one simply has to supply as additional arguments `regformula` and `regdata`. The first one is a formula of the form `~ terms` describing the regression model, and the second is a data frame of regressors. That is, arguments `regformula` and `regdata` are in every respect equivalent to arguments `formula` and `data` of `lm`, with the minor difference that `regformula` does not need to have a left hand side (and is ignored if present). Below, we fit the model

$$X_{it} = \beta_0 + \beta_1 t + \varepsilon_t, \quad t = 1, \dots, 12$$

to the original data set of [Hachemeister \(1975\)](#).

```
> fit <- cm(~state, hachemeister, regformula = ~ time,
+   regdata = data.frame(time = 1:12),
+   ratios = ratio.1:ratio.12,
+   weights = weight.1:weight.12)
> fit
```

```

Call:
cm(formula = ~state, data = hachemeister,
   ratios = ratio.1:ratio.12,
   weights = weight.1:weight.12,
   regformula = ~time, regdata = data.frame(time = 1:12))

Structure Parameters Estimators

Collective premium: 1469 32.05

Between state variance:  24154 2700.0
                        2700  301.8
Within state variance: 49870187

```

To compute the credibility premiums, one has to provide the “future” values of the regressors as in `predict.lm`.

```

> predict(fit, newdata = data.frame(time = 13))
[1] 2437 1651 2073 1507 1759

```

It is well known that the basic regression model has a major drawback: there is no guarantee that the credibility regression line will lie between the collective and individual ones. This may lead to grossly inadequate premiums, as Figure 1 shows.

The solution proposed by [Bühlmann and Gisler \(1997\)](#) is simply to position the intercept not at time origin, but instead at the barycenter of time (see also [Bühlmann and Gisler, 2005](#), Section 8.4). In mathematical terms, this essentially amounts to using an orthogonal design matrix. By setting the argument `adj.intercept` to `TRUE` in the call, `cm` will automatically fit the credibility regression model with the intercept at the barycenter of time. The resulting regression coefficients have little meaning, but the predictions are sensible.

```

> fit2 <- cm(~state, hachemeister, regformula = ~ time,
+           regdata = data.frame(time = 1:12),
+           adj.intercept = TRUE,
+           ratios = ratio.1:ratio.12,
+           weights = weight.1:weight.12)
> summary(fit2, newdata = data.frame(time = 13))

Call:
cm(formula = ~state, data = hachemeister,
   ratios = ratio.1:ratio.12,

```




Figure 1: Collective, individual and credibility regression lines for State 4 of the Hachemeister data set. The point indicates the credibility premium.

```
weights = weight.1:weight.12,
regformula = ~time, regdata = data.frame(time = 1:12),
adj.intercept = TRUE)
```

Structure Parameters Estimators

Collective premium: -1675 117.1

Between state variance: 93783 0
0 8046

Within state variance: 49870187

Detailed premiums

state	Indiv. coef.	Cred. matrix	Adj. coef.
1	-2062.46	0.9947 0.0000	-2060.41
	216.97	0.0000 0.9413	211.10
2	-1509.28	0.9740 0.0000	-1513.59
	59.60	0.0000 0.7630	73.23
3	-1813.41	0.9627 0.0000	-1808.25
	150.60	0.0000 0.6885	140.16
4	-1356.75	0.8865 0.0000	-1392.88
	96.70	0.0000 0.4080	108.77
5	-1598.79	0.9855 0.0000	-1599.89
	41.29	0.0000 0.8559	52.22
Cred. premium			
2457			
1651			
2071			
1597			
1698			

Figure 2 shows the beneficial effect of the intercept adjustment on the premium of State 4.

6 Linear Bayes model

In the pure bayesian approach to the ratemaking problem, we assume that the observations X_t , $t = 1, \dots, n$, of an entity depend on its risk level θ , and that this risk level is a realization of an unobservable random variable Θ . The best (in the mean square sense) approximation to the unknown risk premium $\mu(\theta) = E[X_t|\Theta = \theta]$ based on observations X_1, \dots, X_n is the Bayesian premium

$$B_{n+1} = E[\mu(\Theta)|X_1, \dots, X_n].$$

It is then well known (Bühlmann and Gisler, 2005; Klugman et al., 2012) that for some combinaisons of distributions, the Bayesian premium is linear and can written as a credibility premium

$$B_{n+1} = z\bar{X} + (1 - z)m,$$




Figure 2: Collective, individual and credibility regression lines for State 4 of the Hachemeister data set when the intercept is positioned at the barycenter of time. The point indicates the credibility premium.

where $m = E[\mu(\Theta)]$ and $z = n/(n + K)$ for some constant K .

The combinations of distributions yielding a linear Bayes premium involve members of the univariate exponential family for the distribution of $X|\Theta = \theta$ and their natural conjugate for the distribution of Θ :

- $X|\Theta = \theta \sim \text{Poisson}(\theta)$, $\Theta \sim \text{Gamma}(\alpha, \lambda)$;
- $X|\Theta = \theta \sim \text{Exponential}(\theta)$, $\Theta \sim \text{Gamma}(\alpha, \lambda)$;
- $X|\Theta = \theta \sim \text{Normal}(\theta, \sigma_2^2)$, $\Theta \sim \text{Normal}(\mu, \sigma_1^2)$;
- $X|\Theta = \theta \sim \text{Bernoulli}(\theta)$, $\Theta \sim \text{Beta}(a, b)$;
- $X|\Theta = \theta \sim \text{Geometric}(\theta)$, $\Theta \sim \text{Beta}(a, b)$;

and the convolutions

- $X|\Theta = \theta \sim \text{Gamma}(\tau, \theta)$, $\Theta \sim \text{Gamma}(\alpha, \lambda)$;
- $X|\Theta = \theta \sim \text{Binomial}(\nu, \theta)$, $\Theta \sim \text{Beta}(a, b)$;
- $X|\Theta = \theta \sim \text{Negative Binomial}(r, \theta)$ and $\Theta \sim \text{Beta}(a, b)$.

[Appendix A](#) provides the complete formulas for the above combinations of distributions.

In addition, [Bühlmann and Gisler \(2005, section 2.6\)](#) show that if $X|\Theta = \theta \sim \text{Single Parameter Pareto}(\theta, x_0)$ and $\Theta \sim \text{Gamma}(\alpha, \lambda)$, then the Bayesian estimator of parameter θ — not of the risk premium! — is

$$\hat{\Theta} = \eta \hat{\theta}^{\text{MLE}} + (1 - \eta) \frac{\alpha}{\lambda},$$

where

$$\hat{\theta}^{\text{MLE}} = \frac{n}{\sum_{i=1}^n \ln(X_i/x_0)}$$

is the maximum likelihood estimator of θ and

$$\eta = \frac{\sum_{i=1}^n \ln(X_i/x_0)}{\lambda + \sum_{i=1}^n \ln(X_i/x_0)}$$

is a weight not restricted to $(0, 1)$. (See the "distributions" package vignette for details on the Single Parameter Pareto distribution.)

When argument `formula` is "bayes", function `cm` computes pure Bayesian premiums — or estimator in the Pareto/Gamma case — for the combinations of distributions above. We identify which by means of argument `likelihood` that must be one of "poisson", "exponential", "gamma", "normal", "bernoulli", "binomial", "geometric", "negative binomial" or "pareto". The parameters of the distribution of $X|\Theta = \theta$, if any, and those of the distribution of Θ are specified using the argument names (and default values) of `dgamma`, `dnorm`, `dbeta`, `dbinom`, `dnbinom` or `dpareto1`, as appropriate.

Consider the case where

$$\begin{aligned} X|\Theta = \theta &\sim \text{Poisson}(\theta) \\ \Theta &\sim \text{Gamma}(\alpha, \lambda). \end{aligned}$$

The posterior distribution of Θ is

$$\Theta|X_1, \dots, X_n \sim \text{Gamma}\left(\alpha + \sum_{t=1}^n X_t, \lambda + n\right).$$

Therefore, the Bayesian premium is

$$\begin{aligned}
 B_{n+1} &= E[\mu(\Theta)|X_1, \dots, X_n] \\
 &= E[\Theta|X_1, \dots, X_n] \\
 &= \frac{\alpha + \sum_{t=1}^n X_t}{\lambda + n} \\
 &= \frac{n}{n + \lambda} \bar{X} + \frac{\lambda}{n + \lambda} \frac{\alpha}{\lambda} \\
 &= z\bar{X} + (1 - z)m,
 \end{aligned}$$

with $m = E[\mu(\Theta)] = E[\Theta] = \alpha/\lambda$ and

$$z = \frac{n}{n + K}, \quad K = \lambda.$$

One may easily check that if $\alpha = \lambda = 3$ and $X_1 = 5, X_2 = 3, X_3 = 0, X_4 = 1, X_5 = 1$, then $B_6 = 1.625$. We obtain the same result using `cm`.

```

> x <- c(5, 3, 0, 1, 1)
> fit <- cm("bayes", x, likelihood = "poisson",
+          shape = 3, rate = 3)
> fit
Call:
cm(formula = "bayes", data = x,
   likelihood = "poisson", shape = 3,
   rate = 3)

Structure Parameters Estimators

Collective premium: 1

Between variance: 0.3333
Within variance: 1
> predict(fit)
[1] 1.625
> summary(fit)
Call:
cm(formula = "bayes", data = x,
   likelihood = "poisson", shape = 3,
   rate = 3)

```

Structure Parameters Estimators

Collective premium: 1

Between variance: 0.3333

Within variance: 1

Detailed premiums

Indiv.	mean	Weight	Cred. factor	Bayes premium
2		5	0.625	1.625

A Linear Bayes formulas

This appendix provides the main linear Bayes credibility results for combinations of a likelihood function member of the univariate exponential family with its natural conjugate. For each combination, we provide, other than the names of the distributions of $X|\Theta = \theta$ and Θ :

- the posterior distribution $\Theta|X_1 = x_1, \dots, X_n = x_n$, always of the same type as the prior, only with updated parameters;
- the risk premium $\mu(\theta) = E[X|\Theta = \theta]$;
- the collective premium $m = E[\mu(\Theta)]$;
- the Bayesian premium $B_{n+1} = E[\mu(\Theta)|X_1, \dots, X_n]$, always equal to the collective premium evaluated at the parameters of the posterior distribution;
- the credibility factor when the Bayesian premium is expressed as a credibility premium.

A.1 Bernoulli/beta case

$X|\Theta = \theta \sim \text{Bernoulli}(\theta)$

$\Theta \sim \text{Beta}(a, b)$

$$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Beta}(\tilde{a}, \tilde{b})$$

$$\tilde{a} = a + \sum_{t=1}^n x_t$$

$$\tilde{b} = b + n - \sum_{t=1}^n x_t$$

Risk premium

$$\mu(\theta) = \theta$$

Collective premium

$$m = \frac{a}{a+b}$$

Bayesian premium

$$B_{n+1} = \frac{a + \sum_{t=1}^n X_t}{a + b + n}$$

Credibility factor

$$z = \frac{n}{n + a + b}$$

A.2 Binomial/beta case

$$X|\Theta = \theta \sim \text{Binomial}(\nu, \theta)$$

$$\Theta \sim \text{Beta}(a, b)$$

$$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Beta}(\tilde{a}, \tilde{b})$$

$$\tilde{a} = a + \sum_{t=1}^n x_t$$

$$\tilde{b} = b + n\nu - \sum_{t=1}^n x_t$$

Risk premium

$$\mu(\theta) = \nu\theta$$

Collective premium

$$m = \frac{\nu a}{a+b}$$

Bayesian premium

$$B_{n+1} = \frac{\nu(a + \sum_{t=1}^n X_t)}{a + b + n\nu}$$

Credibility factor

$$z = \frac{n}{n + (a+b)/\nu}$$

A.3 Geometric/Beta case

$$X|\Theta = \theta \sim \text{Geometric}(\theta)$$

$$\Theta \sim \text{Beta}(a, b)$$

$$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Beta}(\tilde{a}, \tilde{b})$$

$$\tilde{a} = a + n$$

$$\tilde{b} = b + \sum_{t=1}^n x_t$$

Risk premium

$$\mu(\theta) = \frac{1 - \theta}{\theta}$$

Collective premium

$$m = \frac{b}{a - 1}$$

Bayesian premium

$$B_{n+1} = \frac{b + \sum_{t=1}^n X_t}{a + n - 1}$$

Credibility factor

$$z = \frac{n}{n + a - 1}$$

A.4 Negative binomial/Beta case

$$X|\Theta = \theta \sim \text{Negative binomial}(r, \theta)$$

$$\Theta \sim \text{Beta}(a, b)$$

$$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Beta}(\tilde{a}, \tilde{b})$$

$$\tilde{a} = a + nr$$

$$\tilde{b} = b + \sum_{t=1}^n x_t$$

Risk premium

$$\mu(\theta) = \frac{r(1 - \theta)}{\theta}$$

Collective premium

$$m = \frac{rb}{a - 1}$$

Bayesian premium

$$B_{n+1} = \frac{r(b + \sum_{t=1}^n X_t)}{a + nr - 1}$$

Credibility factor

$$z = \frac{n}{n + (a - 1)/r}$$

A.5 Poisson/Gamma case

$X|\Theta = \theta \sim \text{Poisson}(\theta)$

$\Theta \sim \text{Gamma}(\alpha, \lambda)$

$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Gamma}(\tilde{\alpha}, \tilde{\lambda})$

$$\tilde{\alpha} = \alpha + \sum_{t=1}^n x_t$$

$$\tilde{\lambda} = \lambda + n$$

Risk premium

$$\mu(\theta) = \theta$$

Collective premium

$$m = \frac{\alpha}{\lambda}$$

Bayesian premium

$$B_{n+1} = \frac{\alpha + \sum_{t=1}^n X_t}{\lambda + n}$$

Credibility factor

$$z = \frac{n}{n + \lambda}$$

A.6 Exponential/Gamma case

$X|\Theta = \theta \sim \text{Exponential}(\theta)$

$\Theta \sim \text{Gamma}(\alpha, \lambda)$

$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Gamma}(\tilde{\alpha}, \tilde{\lambda})$

$$\tilde{\alpha} = \alpha + n$$

$$\tilde{\lambda} = \lambda + \sum_{t=1}^n x_t$$

Risk premium

$$\mu(\theta) = \frac{1}{\theta}$$

Collective premium

$$m = \frac{\lambda}{\alpha - 1}$$

Bayesian premium

$$B_{n+1} = \frac{\lambda + \sum_{t=1}^n X_t}{\alpha + n - 1}$$

Credibility factor

$$z = \frac{n}{n + \alpha - 1}$$

A.7 Gamma/Gamma case

$X|\Theta = \theta \sim \text{Gamma}(\tau, \theta)$

$\Theta \sim \text{Gamma}(\alpha, \lambda)$

$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Gamma}(\tilde{\alpha}, \tilde{\lambda})$

$$\tilde{\alpha} = \alpha + n\tau$$

$$\tilde{\lambda} = \lambda + \sum_{t=1}^n x_t$$

Risk premium

$$\mu(\theta) = \frac{\tau}{\theta}$$

Collective premium

$$m = \frac{\tau\lambda}{\alpha - 1}$$

Bayesian premium

$$B_{n+1} = \frac{\tau(\lambda + \sum_{t=1}^n X_t)}{\alpha + n\tau - 1}$$

Credibility factor

$$z = \frac{n}{n + (\alpha - 1)/\tau}$$

A.8 Normal/Normal case

$X|\Theta = \theta \sim \text{Normal}(\theta, \sigma_2^2)$

$\Theta \sim \text{Normal}(\mu, \sigma_1^2)$

$\Theta|X_1 = x_1, \dots, X_n = x_n \sim \text{Normal}(\tilde{\mu}, \tilde{\sigma}_1^2)$

$$\tilde{\mu} = \frac{\sigma_1^2 \sum_{t=1}^n x_t + \sigma_2^2 \mu}{n\sigma_1^2 + \sigma_2^2}$$

$$\tilde{\sigma}_1^2 = \frac{\sigma_1^2 \sigma_2^2}{n\sigma_1^2 + \sigma_2^2}$$

Risk premium

$$\mu(\theta) = \theta$$

Collective premium

$$m = \mu$$

Bayesian premium

$$B_{n+1} = \frac{\sigma_1^2 \sum_{t=1}^n X_t + \sigma_2^2 \mu}{n\sigma_1^2 + \sigma_2^2}$$

Credibility factor

$$z = \frac{n}{n + \sigma_2^2/\sigma_1^2}$$

References

- H. Belhadj, V. Goulet, and T. Ouellet. On parameter estimation in hierarchical credibility. *ASTIN Bulletin*, 39(2), 2009.
- H. Bühlmann. Experience rating and credibility. *ASTIN Bulletin*, 5:157–165, 1969.
- H. Bühlmann and A. Gisler. Credibility in the regression case revisited. *ASTIN Bulletin*, 27:83–98, 1997.
- H. Bühlmann and A. Gisler. *A course in credibility theory and its applications*. Springer, 2005. ISBN 3-5402575-3-5.
- H. Bühlmann and W. S. Jewell. Hierarchical credibility revisited. *Bulletin of the Swiss Association of Actuaries*, 87:35–54, 1987.
- H. Bühlmann and E. Straub. Glaubwürdigkeit für Schadensätze. *Bulletin of the Swiss Association of Actuaries*, 70:111–133, 1970.
- M. J. Goovaerts and W. J. Hoogstad. *Credibility theory*. Number 4 in Surveys of actuarial studies. Nationale-Nederlanden N.V., Netherlands, 1987.
- V. Goulet. Principles and application of credibility theory. *Journal of Actuarial Practice*, 6:5–62, 1998.
- C. A. Hachemeister. Credibility for regression models with application to trend. In *Credibility, theory and applications*, Proceedings of the Berkeley Actuarial Research Conference on Credibility, pages 129–163, New York, 1975. Academic Press.

- W. S. Jewell. The use of collateral data in credibility theory: a hierarchical model. *Giornale dell'Istituto Italiano degli Attuari*, 38:1–16, 1975.
- S. A. Klugman, H. H. Panjer, and G. Willmot. *Loss Models: From Data to Decisions*. Wiley, New York, 4 edition, 2012. ISBN 978-1-118-31532-3.
- E. Ohlsson. Simplified estimation of structure parameters in hierarchical credibility. Presented at the Zurich ASTIN Colloquium, 2005. URL <https://www.actuaries.org/ASTIN/Colloquia/Zurich/Ohlsson.pdf>.